

Drug Information Sheet("Kusuri-no-Shiori")

External

Published: 09/2015

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: TARIZART OTIC SOLUTION 0.3%

Active ingredient: Ofloxacin

Dosage form: faint yellow to pale yellow limpid aqueous otic solution

Print on wrapping:



Effects of this medicine

This medicine is a new quinolone otic solution. It inhibits DNA replication of bacteria and shows antibacterial effect. It is usually used to treat otitis externa and otitis media.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is<< to be written by a healthcare professional>>
- In general, for adults, apply 6 to 10 drops into the ear canal at a time, twice a day. After applying, a 10-minute ear instillation (keeping the ear facing up) is required. The dosage may be adjusted according to your symptoms. For children, the dosage will be decreased accordingly. Strictly follow the instructions.
- The medicine should be used only for ear dropping.
- Warm the medicine to room temperature before use. If the temperature of the medicine is low, dizziness may occur when applying. Lie down with the ear to be treated facing up when applying the ear drops, and pay attention that the tip of the container does not touch your ear directly. After placing clean gauze on your ear, get up and wipe off any excess liquid.
- If you miss a dose, apply the missed dose as soon as possible. However, if it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. You should not apply two doses at one time.
- If you accidentally apply more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop applying this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include rash, itch, ear pain and headache. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.
- To avoid improper use (misuse) and quality loss (change), do not transfer this solution to any other bottle or container.
- Do not put any other solution in this bottle.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.